

## Identifying Markings

- $\frac{1}{2}$  -  $\frac{3}{4}$  inch long
- Dark smoky brown wings
- One cream/yellow band on second segment of abdomen (near “waist”)
- Three large cream/yellow spots on face



## How to Identify Nests

- Round holes the diameter of a pencil
- Holes go straight down (not angled into the ground)
- Surrounded by a circle of excavated soil (not all to one side like a dog would dig)
- Often tucked beside or partially under a clump of grass
- Nests may be clustered together in a colony. Colonies may contain 5-500 nests.



In New England  
*Cerceris* can be found only in  
**July and August**

(Cerceris is pronounced: 'ser-ser-iss')

(Remember, if you want to monitor a colony for EAB, it should be within 1000 yards of **ash trees**.)

## Biosurveillance

- Carry out biosurveillance at your site 3-4 warm sunny afternoons during July.
- Choose 30-100 nests in one area (as many as you can keep an eye on).
- At each nest, pin the ‘collar’ to the ground with golf tee, placing the second hole over the nest entrance.
- Wasps without prey can come and go easily through the hole (A). A wasp carrying prey will not fit through the hole (B).
- For 1-3 hours, watch as wasps return to nests. Wasps with prey can be netted in flight or caught as they try to pass through the collar. Take prey and release wasp.
- Collect a total of 50 beetles over 3-4 visits. Place beetles from each day in a baggie labeled with place and date. Place in freezer and mail to us at end of summer.




A - Wasp Entering Nest Through Collar

B - Wasp Unable to Enter With Prey  
(photos by P. Careless & M. Bohne)

## Promising Nest Sites

## What is Emerald Ash Borer?

- Hard packed sandy soil
- Areas of human disturbance (**baseball diamonds**, old sand pits, trail and road edges, informal parking lots, fire-pits, etc.)
- Full sunshine
- Sparse vegetation (about 50% hard-packed soil and 50% short vegetation)
- Near a wooded area (~ 200 yards or less)



Typical *Cerceris* Colony Sites  
(photos by P. Careless & C. Teerling)

## Wasp Watcher

- Small metallic green beetle (1/2" long, 1/8" wide)
- An exotic beetle from Asia
- Larva tunnels under the bark
- Attacks and kills **all** species of ash
- First found in Michigan in 2002
- Spreading VERY rapidly across the USA and Canada (primarily in firewood)
- Early detection is difficult. This wasp and WaspWatchers can help.



Female *Cerceris fumipennis*  
with Emerald Ash Borer

For more information, visit our website at  
[www.maine.gov/cerceris](http://www.maine.gov/cerceris)

or contact:

Colleen Teerling  
[colleen.teerling@maine.gov](mailto:colleen.teerling@maine.gov)  
ph: 207 287-3096



*Cerceris fumipennis* is a solitary ground-nesting wasp. The female stocks her nest with buprestid beetles, including emerald ash borer (EAB) when present.

**Biosurveillance** (observing colonies of these native wasps and collecting some of the prey they bring back) is currently the most promising way to monitor for EAB. The Maine Forest Service is looking for colonies of these wasps throughout the state, and would like your help.